

Double-humped Super-luminous Supernovae

D. Leahy, R. Ouyed

University of Calgary, 2500 University Dr NW, Calgary, AB, T2N1N4, Canada

Super-luminous supernova (SLSN) are supernovae showing extreme properties in their light-curves: high peak luminosities (more than 10 times brighter than bright SN Ia), and long durations. Several mechanisms have been proposed for SLSN, such as pair instability SN of a massive progenitor, interaction of the ejecta with a massive circumstellar shell, and the dual-shock quark nova (dsQN) model. The dual-shock quark nova model is unique in that it predicts a normal SN event will be seen ~ 10 days prior to the main SLSN event. The dsQN model is described here and shown that it is consistent with the light curve of the one currently known double-humped SLSN, 2006oz.

1. SUPER-LUMINOUS SUPERNOVAE

Super-luminous supernovae (SLSNe) have been discovered in significant numbers over the past decade. They have high peak brightness, with peak absolute magnitudes of -21 or brighter, and usually have long durations (up to 1 year) compared to normal type I or type II SN. Mechanisms proposed to produce such high luminosities for so long include: pair instability SN (which requires a massive progenitor, $> 100M_{\odot}$); the interaction of fast moving SN ejecta ($\sim 10M_{\odot}$) with a massive circumstellar shell (also with $\sim 10M_{\odot}$); and the dual-shock quark nova (dsQN) model. The quark nova (QN) was proposed as an explanation for SN 2006gy and other SLSNe including SN 2005ap Leahy & Ouyed [2008]. In Ouyed et al. [2009a], we emphasize that the lightcurve of the preceding SN gives a double-humped lightcurve.

In the dsQN model (see section 2 below for details), a normal core-collapse SN explodes to produce a high-mass neutron star. The neutron star converts to a quark star, with a delay of several days, in a violent explosion called a quark nova (QN). The shock produced by the QN then reheats the SN ejecta, which can radiate at high levels for extended periods of time- because the SN ejecta has already expanded so that adiabatic expansion losses are much slower than for the case of a normal SN. It is seen that the dual-shock quark nova model has unique signature: it predicts that a normal core-collapse SN event will be seen several days prior to the main SLSN event.

There is currently one known double-humped SLSN, namely 2006oz Leloudas et al. [2012]. Supernova (SN) 2006oz is a newly-recognized member of the class of H-poor, superluminous supernovae Quimby et al. [2011]. The bolometric light curve shows a precursor event with a duration between 6-10 days in the rest-frame, followed by a dip, after which the luminosity begins to rise. The subsequent rise has previously been fit using three different models: (i) input from radioactive decay; (ii) a magnetar spin-down model; (iii) a circum-stellar medium (CSM) interaction. The Nickel decay model has problems because it requires an unreasonably large

amount ($10.8M_{\odot}$) of ^{56}Ni with a total ejecta mass of $14.4M_{\odot}$. Another problem is that the SN was not detected 9 months later, which is inconsistent with the standard decay curve for ^{60}Co . The magnetar and CSM lightcurve models were shown in Figure 7 of Leloudas et al. [2012]. None of these three models accounts for the precursor event. Yet the dsQN model for SLSN, predicts the existence of a precursor SN Ouyed et al. [2009a] very similar to the observed precursor of SN2006oz.

To show the precursor of SN2006oz is plausibly a normal SN, we estimate its energy be $\sim 10^{49}$ erg $\times t_{\text{pre},10}$ where $t_{\text{pre},10}$ is the duration of the precursor in units of 10 days (limited by the observations from about 7 days to 12 days). This energy is typical of brighter Type-II SNe (e.g. Young [2004]).

In this paper we describe the dual-shock QN (dsQN) model, including the precursor SN. Then we show that the main peak and the precursor of SN2006oz are self-consistently fit by the dsQN model, and conclude with remarks including future work to be done.

2. THE DUAL-SHOCK QUARK NOVA (dsQN) MODEL

A quark nova (QN) was proposed as an alternative explanation for SN 2006gy Leahy & Ouyed [2008], Ouyed et al. [2009a]. A QN is expected to occur when the core density of a neutron star reaches the quark deconfinement density and triggers a violent Ouyed et al. [2002] conversion to the more stable strange quark matter Itoh [1970], Bodmer [1971], Witten [1984]. During the spin-down evolution of the neutron star, accompanied by increasing central density, a detonative Niebergal et al. [2010], Ouyed et al. [2011] phase transition to up-down-strange triplets would result in ejection of the outer heavy element-rich and neutron-rich layers of the neutron star at ultra-relativistic velocities Keranen et al. [2005], Ouyed & Leahy [2009] (in Ouyed & Leahy [2009], see the first panel of Fig. 2). Follow-up studies of neutrino and photon emission processes during the QN Vogt et al.

[2004], Ouyed et al. [2009b] have shown that these outermost layers (of $\sim 10^{-4}$ - $10^{-3}M_{\odot}$ in mass) can be ejected with up to 10^{53} erg in kinetic energy. Nucleo-synthesis simulations of the evolution of the neutron-rich QN ejecta were found to produce primarily heavy elements with mass number, $A > 130$ Jaikumar et al. [2007].

If the time delay (t_{delay}) between SN and QN explosions is too long the SN ejecta will have dissipated such that the QN essentially erupts in isolation. However, when t_{delay} is on the order of days to weeks a violent collision occurs reheating the extended SN ejecta Leahy & Ouyed [2008], Ouyed et al. [2009a]. The brilliant radiance of the re-shocked SN ejecta fades as the photosphere recedes, eventually revealing a mixture of the inner SN ejecta and the QN ejecta material.

3. A PLAUSIBLE CANDIDATE DOUBLE-HUMPED dsQN EVENT: SN2006oz

SN2006oz is the first known double-humped SLSN event Leloudas et al. [2012]. Ouyed & Leahy [2012] study this event in some detail. Here we use SN2006oz as an example of double-humped SNSNe, and argue that, in the dsQN model, we expect to see other similar examples in future.

Figure 1 (from Ouyed & Leahy [2012]) shows the observed SN2006oz light curve using the data from Leloudas et al. [2012]. The g-band data is used, which has the best time coverage and smallest errors for most times. Time is in days at the source using the known redshift ($z = 0.376$). Apparent g-band magnitudes were converted to absolute g-band magnitudes using the corresponding luminosity distance for the standard model Wright [2006]. The suggested extinction correction (B-V) from Leloudas et al. [2012] was included, even though it was small. The dsQN model is also shown in Fig. 1. For the SN lightcurve (the first hump), we use an observed light curve: the light curve of SN1999em from Bersten & Hamuy [2009] which has good time coverage in the first 50 days. Bersten et al. [2011] fitted hydrodynamic models to SN1999em and derived a progenitor mass of $19M_{\odot}$ which is similar in mass to the SN progenitor we used in our QN model. Other parameters for the SN1999em model were progenitor radius of $800R_{\odot}$, explosion energy of 1.25×10^{51} erg and ^{56}Ni mass of $0.056M_{\odot}$. We scaled the bolometric magnitude by +2 to represent a more energetic SN, which is reasonable since the range in brightness of Type II SNe varies considerably with many models giving brighter SN than 1993em (e.g. Young [2004]).

In the QN model the progenitor initial mass is in the range of 20 - $40M_{\odot}$ (see Leahy & Ouyed [2008], Ouyed et al. [2009b], Ouyed et al. [2010])

to create a massive neutron star with core density near the instability to convert to quark matter Niebergal et al. [2010]. This motivates our choice of SN ejected mass of $20M_{\odot}$. Best fits from our previous studies of SLSNe yielded time delays of ~ 10 days which motivates the time delays that we explored. For SN2006oz the shown fit (see Figure 1) uses $t_{\text{delay}} = 6.5$ days, $v_{\text{QN}} = 5000 \text{ km s}^{-1}$ and a preceding SN ejecta with an average velocity of $v_{\text{SN}} \simeq 1900 \text{ km s}^{-1}$. The combined light from the SN and from the QN-reheated SN ejecta give a reasonable fit to the observations with a self-consistent model.

4. COMMENTS

Leloudas et al. [2012] notes the intriguing possibility of an intrinsic precursor event in SN 2005ap-like objects. In the dsQN model, there must be a normal SN ($20 < \text{Mbol} < 15$) preceding the SLSN (if the delay is long enough, $> \sim 10$ days) which should be detectable for nearby SN 2005ap-like explosions. For short delays, the normal SN lightcurve would overlap with the brighter dsQN lightcurve and not give a distinct hump.

SN 2005ap-like events are rare: they occur at a rate of less than one in 10^4 core-collapse SNe Quimby et al. [2011]. dsQNe are also expected to be rare events: the QNe rate is estimated to be $\sim$ one in 1000 core-collapse events with one tenth of them having time delays in the appropriate range to produce dsQNe ($t_{\text{delay}} \sim 5$ - 30 days (Staff et al. [2006], Jaikumar et al. [2007], Leahy & Ouyed [2008], Leahy & Ouyed [2009], Ouyed et al. [2009a]). These two order of magnitude estimates of dsQN events and SN 2005ap-like events are consistent with each other.

We note that the dsQN model applies to both H-rich and H-poor SLSNe, but these occur at similar rates. For both cases, the QN shock reheats the SN envelope to high temperature, so H-poor/H-rich progenitors would give H poor/ H-rich spectra. Because of mass-loss dependency on metallicity, we expect H poor SLSNe to occur in higher-metallicity environments. Low-metallicity progenitors would more likely be H-rich and should in principle have more massive envelopes.

We expect a number of SLSNe to have a double-humped character, as predicted by the dsQN model. The first hump is much fainter and has the brightness of a normal core-collapse SN, but should be observable in relatively nearby SLSNe. It is an for the dsQN model to find additional double-humped SLSNe beyond SN2006oz. We can model the precursor light-curves to learn about the progenitors of SLSNe and of dsQN. The precursor SN is also a clear and unique signature of the dsQN model. Other properties predicted by the dsQN model include: the presence of




Figure 1: (reproduced from Ouyed & Leahy [2012]) SN2006oz r-band lightcurve (solid circles; upper limits shown as triangles). The dsQN model is calculated for $M_{\text{ejecta}} = 20M_{\odot}$ and $(t_{\text{delay}}) = 6.5$ days.

heavy elements Jaikumar et al. [2007], and of spallation nuclei produced in the collision between the fast-moving QN ejecta and the inner parts of the SN ejecta Ouyed et al. [2011].

Acknowledgments

This research is supported the National Science and Engineering Research Council of Canada (NSERC).

References

- Bersten, M. C., & Hamuy, M. 2009, *ApJ*, 701, 200
 Bersten, M. C., Benvenuto, O., & Hamuy, M. 2011, *ApJ*, 729, 61
 Bodmer, A. R. 1971, *Phys. Rev. D*, 4, 1601
 Itoh, N. 1970, *Prog. Theor. Phys.* 44, 291
 Jaikumar, P., Meyer, B. S., Otsuki, K., & Ouyed, R. 2007, *Astronomy& Astrophysics*, 471, 227
 Keränen, P., Ouyed, R., and Jaikumar, P. 2005, *ApJ*, 618, 485
 Leahy, D., and Ouyed, R. 2008, *Mon. Not. Roy. Ast. Soc.*, 387, 1193
 Leahy, D., & Ouyed, R. 2009, *Advances in Astronomy*, 2009
 Leloudas, G., Chatzopoulos, E., Dilday, B., et al. 2012, *Astronomy& Astrophysics*, 541, 129
 Niebergal, B., Ouyed, R., & Jaikumar, P. 2010b, *Phys. Rev. C* 82, 062801 [arXiv:1008.4806 [nucl-th]]
 Ouyed, R., Dey, J., & Dey, M. 2002, *A&A*, 390, L39
 Ouyed, R., & Leahy, D. 2009, *ApJ*, 696, 562
 Ouyed, R., Leahy, D., & Jaikumar, P. 2009a, Proceedings for "Compact stars in the QCD phase diagram II (CSQCD II)", May 20-24, 2009, KIAA at Peking University, Beijing- P. R. China, [http://vega.bac.pku.edu.cn/rxxu/csqed.htm], [arXiv:0911.5424]
 Ouyed, R., Pudritz, R. E., & Jaikumar, P. 2009b, *ApJ*, 702, 1575
 Ouyed, R., Kostka, M., Koning, N., Leahy, D., & Steffen, W. 2010, *MNRAS*, 423, 1652
 Ouyed, R., Leahy, D., Ouyed, A., & Jaikumar, P. 2011, *Physical Review Letters*, 107, 151103
 Ouyed, R., & Leahy, D. 2012, submitted [arXiv:1202.2400v1]
 Quimby, R. M., et al. 2011, *Nature*, 474, 487
 Staff, J. E., Ouyed, R., & Jaikumar, P. 2006, *ApJ*, 645, L145
 Vogt, C., Rapp, R., & Ouyed, R. 2004, *Nuc. Phys. A*, 735, 543
 Witten, E. 1984, *Phys. Rev. D*, 30, 272
 Wright, E. L. 2006, *The Publications of the Astronomical Society of the Pacific*, 118, 1711
 Young, T. R. 2004, *ApJ*, 617, 1233